

August 1992

MPI-Ph/92-65

Non-Abelian Boltzmann Equation for Mixing and Decoherence[†]

G. Raffelt, G. Sigl and L. Stodolsky

*Max-Planck-Institut für Physik
Postfach 401212, 8000 München 40, Germany*

ABSTRACT

We consider particle oscillations and their damping in second-quantized form. We find that the damping or “decoherence” may be described by a Boltzmann-like collision integral with “non-abelian blocking factors” (fermions). Earlier results are generalized in that the momentum degrees of freedom are included and that the mixing equations become intrinsically non-linear at high densities.

PACS index categories: 05.20.Dd, 03.65.Bz, 14.60.Gh, 97.60.Bw

[†] Based, in part, on work to be submitted by G.S. as a doctoral thesis to the Ludwig-Maximilians-Universität (Munich).

1. *Introduction.*—A series of interesting problems, ranging from the question of “decoherence”, such as for the two states of an optical isomer^{1,2}, spin relaxation in condensed matter³, or the damping of particle oscillations⁴, pose essentially the same question in a variety of contexts⁵. For neutrino oscillations in particular, questions of statistics can arise, as in the neutrino degeneracy during the collapsed phase of a supernova. It then becomes of interest to examine a second-quantized approach to such problems, where giving the correct commutation relations to the fields takes care of “Pauli blocking” and similar effects. In addition it might be hoped that the use of field theoretic language may add some insight into the interesting quantum mechanical issues behind our practical applications. In attempting such a formulation we have found a simple and transparent kinetic equation, which may be thought of as a kind of “non-abelian Boltzmann equation”. Although closely related or partial versions of this equation seem to exist in various connections⁶, we think it useful to present it here in a general and simple form.

An interesting aspect of the equation is that it presents kinetic and “interference” aspects simultaneously. Spin-like problems of the type under discussion depend very much on subtle phase effects. On the other hand the Boltzmann type of kinetic equation seems to be entirely “incoherent” and classical in spirit. Nevertheless, we shall see how a marriage of both aspects arises naturally. This also may be of help in understanding many questions including how the classical world arises from the quantum mechanical one.

2. *Matrix of densities.*—In a previous paper⁷, we gave a field-theoretic approach to mixing problems dealing directly with the fields. However, incoherence or damping was not considered. In the single-particle framework used previously⁵ to study damping, the object of study (for two-state systems) was the quantum mechanical 2×2 density matrix for the two states. The two-state system was viewed as imbedded in a much larger system and the density matrix was found by considering the total wavefunction of the whole system and tracing over the unobserved variables of the large system—usually thought of as “the medium”. If the density matrix was expressed in terms of Pauli matrices $\boldsymbol{\tau}$ and a polarization vector $\mathbf{P}$ as $\frac{1}{2}(1 + \mathbf{P} \cdot \boldsymbol{\tau})$ then $\mathbf{P}$ was found to obey a Bloch-like equation:

$$\dot{\mathbf{P}} = \mathbf{V} \times \mathbf{P} - D\mathbf{P}_T. \quad (1)$$

This gives a coherent rotation of the polarization through $\mathbf{V}$, and through the damp-

ing parameter D , a loss of coherence or damping manifested as a shortening of $\mathbf{P}$. The subscript T indicates those components non-diagonal (“transverse”) with respect to interactions with the medium. Also, a formula for D was obtained in S -matrix language, resembling a generalized optical theorem, and giving D in terms of the scattering amplitudes on the background. In measurement theory language, D may be thought of as giving the rate of “reduction” of the wavefunction of the two-state system⁵.

What is to play the role of the quantum mechanical 2×2 density matrix in the field-theoretic formulation? In second-quantized terms the above procedure of tracing over the unobserved variables may be accomplished in a sense by applying an annihilation operator to the state to “remove” the coordinate and then finding the overlap with another such operation. Thus we propose to study the “matrix of densities”

$$\rho_{ij} = \langle |\psi_j^\dagger \psi_i| \rangle \quad (2)$$

instead of the density matrix. The indices refer to the various components of the field in question and $|\rangle$ to our original state. For a two-state system, such as neutrino fields with two flavors, ψ will be a two-component spinor in flavor space⁸, and we will have four bilinears to deal with. Diagonal elements of this c-number matrix are the ordinary densities of a particle type, while the off-diagonal elements contain more subtle phase information relating to the coherence of the system. The matrix (2) is Hermitian and positive definite. If we assume interactions that conserve the total number of particles, as we shall in this paper, then the trace of its spatial integral is conserved. Hermiticity, positivity and conservation of the trace are the properties characterizing a density matrix and so it is justified to speak of the integrated ρ as a density matrix⁹.

The object (2) is of course much more complicated than the density matrix of the single particle formulation since the arguments of the fields can refer to any, and in general different, points of space; similarly in momentum space they can refer to different momenta. However, since our major interest is the mixing question, we shall make the greatly simplifying assumption of ignoring spatial inhomogeneities. Our state, $|\rangle$, will represent a situation which is spatially uniform. Thus in position space ρ can only depend on the difference of the arguments of the fields, while in momentum space we have $\langle |\psi_j^\dagger(\mathbf{p}') \psi_i(\mathbf{p})| \rangle = (2\pi)^3 \delta^3(\mathbf{p}' - \mathbf{p}) \rho_{ij}(\mathbf{p})$, and so we can confine ourselves to the study of the matrix $\rho_{\mathbf{p}}$. (When matrix indices are suppressed

we write $\mathbf{p}$ as a subscript).

In this paper we shall limit ourselves to the non-relativistic domain in the sense that we do not directly consider the role of creation and annihilation of particles. Thus our fields are not those of fully relativistic quantum field theory; the operator $\psi(\mathbf{x})$ contains only annihilation operators for particles and not the creation operators for antiparticles.

3. Interactions.—Turning to the time development of ρ , we treat the fields as Heisenberg operators $\psi(t)$ and assume that our constant state $|\rangle$ is produced by letting the creation operators for the ψ particles operate on the state vector for the medium at $t = 0$. This corresponds to a simple product wavefunction with no correlations between the ψ -particles and the medium. We follow the approach of finding the equations among the ρ_{ij} by turning on the interactions for a short time at $t = 0$ and then assuming these equations to be valid at all times.

Our starting point is the equation of motion for the operator $\psi_j^\dagger \psi_i$, the Hamiltonian consisting of a “free” and an interaction term, $H = H_{\text{free}} + H_{\text{int}}$. H_{free} is taken to be bilinear in the fields and diagonal in momentum. It may describe a kinetic energy plus some internal interaction of the fields among themselves. An example is the free Hamiltonian for neutrinos with a mass matrix, which leads to a rotation in the internal flavor space. If (i, j) refer to different spin components of a system with a magnetic moment, H_{free} may refer to an applied magnetic field, also leading to a rotation among the components. Or for the molecular isomers H_{free} gives the tunneling between left and right configurations in addition to the kinetic energy.

H_{free} may be written for each momentum in terms of a matrix Ω and the column vector ψ as $\psi_{\mathbf{p}}^\dagger \Omega_{\mathbf{p}} \psi_{\mathbf{p}}$. Commuting with $\psi_j^\dagger(\mathbf{p}) \psi_i(\mathbf{p})$ produces bilinears of the same momentum. The set of densities “rotate” among each other. This is of course just the precession of the “spin”, ordinary or flavor variety.

H_{int} gives the interaction with the background medium which we take to be a local point-like interaction between the ψ and the background, represented by an operator B :

$$H_{\text{int}}(t) = \int B(x) \psi^\dagger(x) G \psi(x) d^3 \mathbf{x}. \quad (3)$$

G is a coupling constant matrix acting on the column vector of the ψ 's, and B would be typically, as in the Fermi interaction, bilinear in the background fields. When G is diagonalized it gives a set of states which scatter only into themselves. The T for “transverse” in Eq. (1) then refers to those bilinears which are off-diagonal in such

states. The choice of Eq. (3) is appropriate to the Fermi interaction and many other problems where the particles under study undergo a local elastic scattering as their basic interaction. We shall take G Hermitian, corresponding to lowest order, but one can imagine relaxing this restriction to represent an underlying process in higher than Born approximation, as in the S -matrix approach^{1,2}.

The commutator of H_{int} with $\psi_j^\dagger(\mathbf{p})\psi_i(\mathbf{p})$ again yields a bilinear. Finally taking the expectation value of the resulting equation of motion yields [notation: $d\mathbf{p} \equiv d^3\mathbf{p}/(2\pi)^3$]:

$$i\dot{\rho}_{ij}(t, \mathbf{p}) = [\Omega(\mathbf{p}), \rho(t, \mathbf{p})]_{ij} + \int d\mathbf{p}' \langle [B(t, \mathbf{p} - \mathbf{p}')\psi_j^\dagger(t, \mathbf{p})G_{ik}\psi_k(t, \mathbf{p}')]\rangle - \text{h.c.} \quad (4)$$

This is an exact expression in terms of Heisenberg operators. It resembles but is not yet in the desired form of a set of equations among the ρ_{ij} , like Eq. (1), because B cannot be factorized out of the brackets. The original state $|\rangle$ is given by free particle operators, $\psi^\dagger$ at $t = 0$, while the operator expressions in the brackets are non-trivial in terms of such Schrödinger operators.

We can, however, introduce perturbation theory where we expand the Heisenberg operators $\psi(t)$ and $B(t)$ in terms of non-interacting (interaction representation) operators $\psi_0(t)$ and $B_0(t)$ corresponding to $G = 0$. A first approximation is to set $\psi(t) = \psi_0(t)$ and $B(t) = B_0(t)$. Now with this approximation the bracket in Eq. (4) may be factorized since by assumption the original state $|\rangle$ has no correlations between the ψ -particles and the background. Only bilinears with equal momenta survive by virtue of the spatial uniformity represented by $|\rangle$ and we obtain a closed equation among the ρ :

$$i\dot{\rho}_{\mathbf{p}} = [\Omega_{\mathbf{p}}, \rho_{\mathbf{p}}] + \rho_{\text{med}} [G, \rho_{\mathbf{p}}] . \quad (5)$$

Here, $\rho_{\text{med}} \equiv \langle |B(0, 0)| \rangle$ is the medium density, assumed to be stationary and taken at momentum transfer zero. This relation will be recognized as adding an “index of refraction” or medium-induced energy contribution to the free mixing given by H_{free} . Note this index of refraction effect actually refers to *differences* in the index (as occur for example as in the MSW effect) and does not appear for a one-component field.

In the two-level [or more generally $\text{SU}(2)$] case where Ω , ρ and G can be expanded in terms of Pauli matrices [more generally $\text{SU}(2)$ generators], *e.g.* $\rho_{\mathbf{p}} = \frac{1}{2}n_{\mathbf{p}}(1 + \mathbf{P}_{\mathbf{p}} \cdot \boldsymbol{\tau})$, with coefficients $\mathbf{V}_{\text{free}}$, $\mathbf{P}$ and $\mathbf{G}$ respectively, Eq. (5) becomes

$\dot{\mathbf{P}}_{\mathbf{p}} = \mathbf{V}_{\mathbf{p}} \times \mathbf{P}_{\mathbf{p}}$ with $\mathbf{V}_{\mathbf{p}} \equiv \mathbf{V}_{\text{free}}(\mathbf{p}) + \mathbf{V}_{\text{med}}$ where $\mathbf{V}_{\text{med}} = \mathbf{G}\rho_{\text{med}}$. This is the first term of Eq. (1) and agrees with the old result⁵, to first order in G .

4. *Incoherence and damping.*—Now, Eq. (5) still contains no sign of any loss of coherence, *i.e.* any shrinking of $\mathbf{P}$. This is understandable, since to this order of approximation there is no excitation of the medium. Only forward scattering has been taken into account, as is indicated by taking B at momentum transfer zero. If the medium does not change its state, no incoherence can be expected since the overall wavefunction, assumed to be a simple product at $t = 0$, has remained so. As is familiar in perturbation theory, the energy but not the wavefunction changes in first order. Thus to see the first evidence of damping or incoherence we must go to the next order.

We stress that spatial homogeneity excludes sources of damping or incoherence due to spatially varying conditions. For example, if the magnetic field applied to an ensemble of spins varies in space, or if there are inhomogeneities in the medium in which our neutrinos propagate¹⁰, there will be an averaging or loss of coherence in the properties of the ensemble. In general a loss of coherence can always result from averaging over some parameters characterizing the system. However, here we only wish to address the true quantum mechanical loss of coherence.

We go to the next order in G by expanding the Heisenberg operators to first order:

$$\begin{aligned}\psi(t) &= \psi_0(t) + i \int_0^t dt' [H_{\text{int}}(t - t'), \psi_0(t)], \\ B(t) &= B_0(t) + i \int_0^t dt' [H_{\text{int}}(t - t'), B_0(t)].\end{aligned}\tag{6}$$

To evaluate the new terms in Eq. (6) we must now make our assumptions on the nature of the system explicit. We assume that the interactions described by H_{int} can be taken as individual, isolated collisions where the particles represented by the ψ go from free states to free states as in ordinary scattering theory. Thus the time for a single collision is taken to be short compared to the time between collisions; furthermore the very small energies represented by the mixing effects in Eq. (5) are neglected in calculating the effects of the collisions. Under these assumptions we can carry out what amounts to a Golden Rule calculation on the level of the fields and

find

$$\begin{aligned}
\psi(t, \mathbf{p}) &= \psi_0(t, \mathbf{p}) + ie^{-iE_{\mathbf{p}}t} \int d\mathbf{p}' d\omega \delta(E_{\mathbf{p}'} - E_{\mathbf{p}} + \omega) B_0(\omega, \mathbf{p} - \mathbf{p}') G \psi_0(0, \mathbf{p}'), \\
B(t, \mathbf{p} - \mathbf{p}') &= B_0(t, \mathbf{p} - \mathbf{p}') + i \int d\mathbf{q}' d\mathbf{q} d\omega \delta(E_{\mathbf{q}'} - E_{\mathbf{q}} - \omega) \times \\
&\quad \times [B_0(\omega, \mathbf{q}' - \mathbf{q}), B_0(t, \mathbf{p} - \mathbf{p}')] \psi_0^\dagger(0, \mathbf{q}') G \psi_0(0, \mathbf{q}).
\end{aligned} \tag{7}$$

The δ function arises from letting the time integrals go to infinity. We have dropped an associated principle-part integral since this has to do with a second order energy shift in the medium, which we already have to lowest order in Eq. (5).

We may now substitute Eq. (7) into the r.h.s. of Eq. (4). By construction the state $|\rangle$ has the property that only bilinear correlations of the type $\langle |\psi^\dagger \psi| \rangle$ are present, so in carrying out the contractions in the brackets only ρ 's will arise. It must be recognized, however, that an assumption is involved here, similar to that involved in the classical Boltzmann equation. While we construct the state at $t = 0$ such that no other correlations are present, the philosophy of the method that the equation found by turning on the interactions for a short time at $t = 0$ can be used for all times tacitly assumes that no other kinds of correlations can build up significantly. In this sense our derivation is Boltzmann-like and, for example, we exclude the possibility that our system might have important pair correlations.

Carrying out, then, the contractions with only the survival of ordinary density bilinears, and evaluating all quantities at $t = 0$, we arrive at our main result:

$$\begin{aligned}
\dot{\rho}_{\mathbf{p}} &= -i [\Omega_{\mathbf{p}}, \rho_{\mathbf{p}}] - i \rho_{\text{med}} [G, \rho_{\mathbf{p}}] \\
&\quad + \int d\mathbf{p}' \left(S(p' - p) \frac{(1 - \rho_{\mathbf{p}}) G \rho_{\mathbf{p}'} G + \text{h.c.}}{2} - S(p - p') \frac{G(1 - \rho_{\mathbf{p}'}) G \rho_{\mathbf{p}} + \text{h.c.}}{2} \right)
\end{aligned} \tag{8}$$

where $S(\Delta)$, a function of the four-vector Δ , is the usual dynamical structure function, defined by $\int_{-\infty}^{+\infty} dt e^{i\Delta_0 t} \langle |B_0(t, \mathbf{\Delta}) B_0(0, -\mathbf{\Delta})| \rangle$. This is the desired generalization of Eq. (1) under Boltzmann-like conditions. It resembles the ordinary Boltzmann equation with the usual drift term, absent because of spatial uniformity, replaced by the commutators for coherent rotations, which give the free motion in “spin space”. The principle new features of Eq. (8) relative to Eq. (1) is the inclusion of the momentum degrees of freedom and the nonlinearity in ρ introduced by statistics. The

main difference with the ordinary Boltzmann equation is the matrix structure of the “collision term”. The “loss” term contains such effects as the creation of polarization by absorption while the “gain” term gives effects like the transfer of polarization through scattering.

We have taken the original state $|\rangle$ to be pure. If it is itself characterized by a density matrix then our various bracketed expressions can be interpreted as thermal or other averages. Similarly, if there are several species of background particles, uncorrelated with one another, their contributions can be summed separately in the collision integral. Eq. (3) is the simplest possible interaction. However, as long as the essential feature that the interactions of the different ψ with the background are identical up to a coupling constant is retained, we anticipate no change from more general interactions except for a generalization of the structure factor S . For example, when the spin structure of the background is accounted for $S(p - p')$ will be replaced¹¹ by the probability $W(p, p')$, which is no longer simply a function of the difference $(p - p')$. It may be verified that this has no effect on our arguments.

By integrating over $\mathbf{p}$ we may also study the total density matrix, with its associated total polarization:

$$\dot{\rho} = -i\rho_{\text{med}}[G, \rho] - i \int [\Omega_{\mathbf{p}}, \rho_{\mathbf{p}}] d\mathbf{p} + \frac{1}{2} \int S(p - p') \left[G, \left(\rho_{\mathbf{p}} G (1 - \rho_{\mathbf{p}'}) - \text{h.c.} \right) \right] d\mathbf{p} d\mathbf{p}'. \quad (9)$$

We see here from the commutator structure that the total number of particles, given by $\text{Tr}(\rho)$, is explicitly conserved. Observe that in the special case where all types scatter with the same amplitude on the medium, *i.e.* G proportional to the unit matrix, the collision term in Eq. (9) vanishes identically. There is then no dynamical damping, however there remain “de-phasing” effects due to the momentum dependence in the precession of the different modes.

5. Limiting cases and applications: Ordinary collision integral.—The simplest limiting case obtains when we have a one-component ψ (or equivalently many components but no mixing terms, or mixing terms which can be simultaneously diagonalized with G) and the background is simply free particles, *i.e.* $B(\mathbf{x}) = \phi^\dagger(\mathbf{x})\phi(\mathbf{x})$, $\phi(\mathbf{x})$ a free field. Then:

$$S(\Delta) = \int d\mathbf{q} d\mathbf{q}' (2\pi)^4 \delta^4(\Delta + q - q') n_{\text{med}}(\mathbf{q}) [1 \pm n_{\text{med}}(\mathbf{q}')], \quad (10)$$

where $n_{\text{med}}(\mathbf{q})$ are the occupation numbers in the medium ($\pm$ for bosons or fermions).

In the low density limit where the blocking factor can be neglected, with a single medium mode $\mathbf{q}_{\text{med}}$, $n(\mathbf{q}) = (2\pi)^3 \rho_{\text{med}} \delta^3(\mathbf{q} - \mathbf{q}_{\text{med}})$, S becomes simply the energy δ function times $2\pi\rho_{\text{med}}$. Because $G^2 \int d\mathbf{p}' S$ is the scattering rate, Eq. (8) has the appearance of standard rate-of-gain and -loss terms. With only a single mode $\mathbf{p}$ of ψ populated, as for a beam, one finds $\dot{\rho}_{\mathbf{p}}/\rho_{\mathbf{p}} = -G^2 \int d\mathbf{p}' S(p - p')$ giving the expected depletion of the beam. S can of course also account for non-trivial interactions among the background particles; it will then have more structure than simply the δ function.

Bloch equations.—An opposite, more “quantum mechanical” limit occurs when S depends negligibly on the energy transfer, as can happen, say for a fixed spin, or more generally when the energy transfer is small compared to the scale of energies associated with processes in the medium. Then $S(\Delta) \approx S(\mathbf{\Delta}) = S(-\mathbf{\Delta})$ and we find that the non-linear terms in Eq. (9) disappear, leaving for the matrix structure of the collision term $2G\rho G - GG\rho - \rho GG = -[G, [G, \rho]]$, which puts the nature of the collision term as a kind of double commutator in evidence. The disappearance of the non-linear terms is interesting; self-blocking by the ψ_i does not reduce the damping. Evidently it should not be thought that the particles do not scatter even if they are not permitted to change their state in some particular basis. It is in this way, after all, that a degenerate Fermi gas exerts a pressure. Blocking effects in the *background* as in Eq. (10), on the other hand, are *not* to be ignored.

Carrying out the integrations in Eq. (9) gives an equation solely in terms of the total ρ (for Ω independent of $\mathbf{p}$), leading to a Bloch-like³ equation. In the two-level case this gives for the collision term:

$$\frac{1}{2} \int d\mathbf{p} d\mathbf{p}' S(\mathbf{p} - \mathbf{p}') n_{\mathbf{p}} \mathbf{G} \times (\mathbf{G} \times \mathbf{P}) = -\frac{1}{2} \int d\mathbf{p} d\mathbf{p}' S(\mathbf{p} - \mathbf{p}') n_{\mathbf{p}} |\mathbf{G}|^2 \mathbf{P}_T = -D \mathbf{P}_T. \quad (11)$$

T means the direction perpendicular to $\mathbf{G}$, where $G = \frac{1}{2}(G_0 + \mathbf{G} \cdot \boldsymbol{\tau})$. This is the damping term of Eq. (1). The calculations given here verify, by essentially standard manipulations, the results of the more intuitive S -matrix arguments, where D was found for two states (“up” and “down”) as the real part of $(S_{\text{up}} S_{\text{down}}^\dagger - 1)$ or in terms of phase shifts to lowest order, $D \sim (\delta_{\text{up}} - \delta_{\text{down}})^2$, which is indeed $\sim (g_{\text{up}} - g_{\text{down}})^2$, as in Eq. (11). This contains, for example, the old result² that if one type of particle interacts and the other not, that the damping parameter is equal to half of the (unblocked) collision rate of the interacting type.

Sign of D and Spontaneous Polarization.— D in Eq. (11) has arrived with the correct sign—we have damping and not anti-damping. This “arrow of time” arises from straightforward calculation—at least for the short time the interaction is on. It is a consequence of our starting from an uncorrelated state, which can only become more correlated in time. That the damping sign is not entirely trivial is revealed by numerical solutions of Eq. (8) where we find when spin and momentum degrees of freedom are closely coupled that the relaxation of the momentum can lead to a transient polarization for an initially unpolarized situation.

Neutrino Mixing in a Stochastic Background.—Flavor mixing in a supernova would be of significance since it allows other neutrinos to share the large ν_e degeneracy. In previous work the approach to kinetic and chemical equilibrium was treated in the single-particle wavefunction approach¹². Our kinetic equation, with degeneracy effects built in, is well suited to this problem. Even though our treatment here is not explicitly relativistic, relativistic kinematics can be incorporated in $\Omega_{\mathbf{p}}$, and the absence of the anti-particle degrees of freedom is not critical since these are suppressed by the high chemical potential. It is important, however, to include $\nu_e - e$ conversions; this may be done by introducing further bilinears involving the electrons, a detailed account will be given elsewhere¹³. For applications where the anti-particles are important, the formalism should be extended, which raises the interesting question of dealing with relativistic fields.

Understanding “Decoherence”.—We recall⁵ that in the first quantized formulation the value of $\mathbf{P}_T$ gives the extent to which the eigenstates of G , “up” and “down” (two-level case), can interfere. As $\mathbf{P}_T$ is reduced the two states “decohere”. Similarly here, interference terms between operators ψ_i corresponding to different eigenvalues of G are given by the appropriate elements of ρ and are generally decreased by the action of the collision term. Hence we may say that in the collision integral we are finding the (rate of) “decoherence”. The damping or decoherence rate is like the scattering rate by the background, but with the difference amplitude $\sim (g_{\text{up}} - g_{\text{down}})$; *difference* because the scattering must distinguish the two systems in order to “measure” and “reduce” them. Hence interactions not distinguishing the states in question are not “reducing”. The other extreme, that of strong damping, leads through the rapid disappearance of $\mathbf{P}_T$, to the “freezing” of the other component¹⁴ and hence to the “Turing-Xeno-Watched-Pot” effect², giving the stabilization¹ of the molecular isomers.

We are grateful to W. Götze for extensive conversations on relaxation theory, to D. Einzel for a discussion concerning spin effects in He^3 and to P. Breitenlohner for a helpful remark.

REFERENCES

1. R. A. Harris and L. Stodolsky, *J. Chem. Phys.* **74** (1981) 2145 and *Phys. Lett.* **78 B** (1978) 313; R. A. Harris and R. Silbey, *J. Chem. Phys.* **78:12** (1983) 7330.
2. R. A. Harris and L. Stodolsky, *Phys. Lett.* **116 B** (1982) 464.
3. A. Abragam, *The Principles of Nuclear Magnetism* (Oxford, 1961), Ch. viii, gives a standard discussion of spin relaxation. Harris and Silbey, Ref. 1, also give numerous references on relaxation theory.
4. For neutrinos see L. Stodolsky *Phys. Rev. D* **36** (1987) 2273, an extensive application of the formalism in the early universe is given by K. Enquist, K. Kainulainen and J. Maalampi, *Nucl. Phys.* **B 349** (1991) 754.
5. A review and discussion is given by L. Stodolsky, in: J. S. Anandan (ed.), *Quantum Coherence* (World Scientific, Singapore, 1990) pg. 320.
6. See for example the discussion of spin effects in He^3 by D. Einzel, in: M. J. R. Hoch and R. H. Lemmer (eds.), *Low Temperature Physics*, (Springer, Heidelberg, 1991) pg. 275.
7. G. Raffelt, G. Sigl, and L. Stodolsky, *Phys. Rev. D* **45** (1992) 1782.
8. A. Dolgov, *Sov. J. Nucl. Phys.* **33** (1981) 700, also used such flavor densities in the description of neutrino mixing.
9. For related attempts to generalize the formalism see M. J. Thomson, *Phys. Rev. A* **45** (1992) 2243; M. J. Thomson and B. H. J. McKellar, University of Melbourne Preprint UMP 89-108 and Manchester preprint MC-TP-92.
10. R. F. Sawyer, *Phys. Rev. D* **42** (1990) 3908.
11. As in N. Iwamoto and C. J. Pethick, *Phys. Rev. D* **25** (1982) 313.

12. J. Maalampi and J. T. Peltoniemi, *Phys. Lett.* **269 B** (1991) 357. M. Turner, *Phys. Rev. D* **45** (1992) 1066.
13. G. Raffelt and G. Sigl, “Neutrino Flavor Conversion in a Supernova Core”, Report MPI-Ph/92-68 (1992).
14. For a discussion of this phenomenon in the $SU(N)$ case see I. B. Khriplovich and V. V. Sokolov, *Physica* **A141** (1987) 73. A further question one could study by the present methods is that of the possibility of interference effects between different degenerate states of large, quasi-macroscopic systems; a big molecule, after all, is almost such a system. If the system has many equivalent states the $SU(N)$ analysis then becomes relevant.